

ORBITAL PHASE IN INSPIRALLING COMPACT BINARIES *

MÁTYÁS VASÚTH[†], BALÁZS MIKÓCZI[‡] and LÁSZLÓ Á. GERGELY[†]

[†]*KFKI Research Institute for Particle and Nuclear Physics
Budapest 114, P.O.Box 49, H-1525, Hungary*

[‡]*Departments of Theoretical and Experimental Physics, University of Szeged
Dóm tér 9, Szeged H-6720, Hungary*

vasuth@rmki.kfki.hu, mikoczi@titan.physx.u-szeged.hu, gergely@physx.u-szeged.hu

We derive the rate of change of the mean motion up to the second post-Newtonian order for inspiralling compact binaries with spin, mass quadrupole and magnetic dipole moments on eccentric orbits. We give this result in terms of orbital elements. We also present the related orbital phase for circular orbits.

Keywords: compact binaries, post-Newtonian expansion, spin, quadrupole moment

Observations by Earth-based gravitational wave observatories are under way aiming to detect gravitational radiation. Upper limits from interferometer data were already set on inspiral event rates for both binary neutron stars¹ and binaries of 3–20 solar mass black holes.² The parameters of spinning compact binaries can be estimated and alternative theories of gravity can also be tested from these measurements.³

An important characteristic of these binaries is the rate of decrease of the orbital period T due to the energy and angular momentum carried away by gravitational waves. Here we give the radiative change of the mean motion $n = 2\pi/T$ (for eccentric orbits). We also present the related change occurred in the orbital phase (for circular orbits). In both expressions we include all known *linear* perturbations for an isolated compact binary. These are the post-Newtonian (PN), spin-orbit (SO), spin-spin (SS), self spin (Self, quadratic in the single spins), quadrupole-monopole (QM) and magnetic dipole-magnetic dipole (DD) contributions.

The expression of the radial period, defined as half of the time elapsed between the turning points, emerges from generic considerations on the perturbed Keplerian motion.^{4,5} Collecting all linear contributions the mean motion has the following form

$$n = \frac{(2\mathcal{E})^{3/2}}{Gm} \left[1 - (15 - \eta) \frac{\mathcal{E}}{4c^2} \right], \quad (1)$$

where $\eta = \mu/m$ is the ratio of the reduced mass μ to the total mass m of the binary system, and $\mathcal{E} = -E/\mu$ where E is the conserved energy. Remarkably there are no explicit spin, quadrupolar and magnetic dipolar contributions in the functional form of the mean motion. These however contribute implicitly to n through $\mathcal{E}$. Since the mean motion is a function of E alone, its evolution can be computed as

* Research supported by OTKA grants nos. T046939, TS044665, F049429 and the János Bolyai Fellowships of the Hungarian Academy of Sciences. M.V. and L.Á.G. wish to thank the organizers of the 11th Marcel Grossmann Meeting for support.

2 *Vasúth, Mikóczy, Gergely*

$\langle dn/dt \rangle = -1/\mu \langle dn/d\mathcal{E} \rangle \langle dE/dt \rangle$. All linear contributions to the secular energy loss $\langle dE/dt \rangle$ due to finite size effects are explicitly given in the literature,⁶⁻⁹ in terms of dynamical constants. The PN contribution is also well-known.¹⁰ Employing these we find the change of the mean motion:

$$\left\langle \frac{dn}{dt} \right\rangle_N = N_0 , \quad (2)$$

$$\left\langle \frac{dn}{dt} \right\rangle_{PN} = - \frac{Gm}{56c^2a(1-e^2)} [N_1 + N_2\eta] , \quad (3)$$

$$\left\langle \frac{dn}{dt} \right\rangle_{SO} = - \frac{G^{1/2}}{2c^2a^{3/2}(1-e^2)^{3/2}m^{1/2}} [N_3S_L + N_4\Sigma_L] , \quad (4)$$

$$\left\langle \frac{dn}{dt} \right\rangle_{Self} = \frac{1}{64c^2a^2(1-e^2)^2} \sum_{i=1}^2 \left(\frac{S_i}{m_i} \right)^2 [N_5 \sin^2 \kappa_i \cos 2(\psi_0 - \psi_i) + N_6(6 + \sin^2 \kappa_i)] , \quad (5)$$

$$\left\langle \frac{dn}{dt} \right\rangle_{SS} = \frac{S_1 S_2}{32c^2 m \mu a^2 (1-e^2)^2} \times [N_7 \sin \kappa_1 \sin \kappa_2 \cos 2(\psi_0 - \bar{\psi}) + N_8 \cos \kappa_1 \cos \kappa_2 + N_9 \cos \gamma] , \quad (6)$$

$$\left\langle \frac{dn}{dt} \right\rangle_{QM} = \frac{m^2}{4a^2(1-e^2)^2} \sum_{i=1}^2 p_i [N_{10}(2 - 3 \sin^2 \kappa_i) + N_{11} \sin^2 \kappa_i \cos 2(\psi_0 - \psi_i)] , \quad (7)$$

$$\left\langle \frac{dn}{dt} \right\rangle_{DD} = - \frac{d_1 d_2}{2Gm\mu a^2(1-e^2)^2} (N_{10}\mathcal{A}_0 + N_{11}\mathcal{B}_0) , \quad (8)$$

where we have introduced the notations $S_L = (S_1 \cos \kappa_1 + S_2 \cos \kappa_2)$, $\Sigma_L = [(m_2/m_1) S_1 \cos \kappa_1 + (m_1/m_2) S_2 \cos \kappa_2]$ and

$$N_j = \frac{G^{7/2} m^{5/2} \mu}{5c^5 a^{11/2} (1-e^2)^{7/2}} \sum_{i=0}^3 n_{ij} e^{2i}, \quad j = 0..11 , \quad (9)$$

Here κ_i and ψ_i are the polar and azimuthal angles of the i^{th} spin vector and the numeric coefficients n_{ij} are:

i	j=0	1	2	3	4	5	6	7	8	9	10	11
0	96	28016	9408	2128	1680	0	64	-3072	-194368	65216	2888	288
1	292	160248	43120	7936	7924	16	608	100112	-621536	211232	9660	-7924
2	37	34650	20916	3510	4224	80	552	113248	-264792	91944	3897	-8570
3	0	-5501	-1036	363	291	9	36	8937	-4500	1740	187	-464

The orbital elements a , e were derived¹¹ from the turning points of the radial part of the perturbed motion cf. $r_{\min}^{max} = a(1 \pm e)$. (In these variables $n = (Gm/a^3)^{1/2} [1 + (\eta - 9) Gm/2ac^2]$).

For Keplerian motions the orbital frequency $\omega = n$. Due to the perturbations, precessions occur in the plane of motion (like periastron advance), and the plane of motion can also evolve. Therefore the relation of ω and n becomes more complicated.¹² In the presence of the PN, SO, SS, Self, QM and DD perturbations,

for circular orbits ($e = 0$) the change in the orbital frequency due to gravitational radiation is:^{13,14}

$$\left\langle \frac{d\omega}{dt} \right\rangle = \frac{96\eta m^{5/3} \omega^{11/3}}{5} \left[1 - \left(\frac{743}{336} + \frac{11}{4}\eta \right) (m\omega)^{2/3} + (4\pi - \beta) m\omega + \left(\frac{34103}{18144} + \frac{13661}{2016}\eta + \frac{59}{18}\eta^2 + \sigma \right) (m\omega)^{4/3} \right], \quad (10)$$

where $\sigma = \sigma_{SS} + \sigma_{Self} + \sigma_{QM} + \sigma_{DD}$ and $\beta, \sigma_{S_1 S_2}, \sigma_{Self}, \sigma_{QM}, \sigma_{DD}$ are the spin-orbit, spin-spin, self-interaction spin, quadrupole-monopole and magnetic dipole-dipole parameters.¹⁴ For completeness we have added the 2PN and tail contributions¹⁵ and we note that higher order contributions are also known.^{16,17}

In terms of the dimensionless time variable $\tau = \eta(t_c - t)/5m$, defined in terms of the time $(t_c - t)$ left until the final coalescence, the accumulated orbital phase is $\phi = \phi_c - (5m/\eta) \int \omega(\tau) d\tau$, where ϕ_c is an integration constant. To second post-Newtonian order:

$$\phi = \phi_c - \frac{1}{\eta} \left\{ \tau^{5/8} + \left(\frac{3715}{8064} + \frac{55}{96}\eta \right) \tau^{3/8} + \frac{3}{4} \left(\frac{\beta}{4} - \pi \right) \tau^{1/4} + \left(\frac{9275495}{14450688} + \frac{284875}{258048}\eta + \frac{1855}{2048}\eta^2 - \frac{15\sigma}{64} \right) \tau^{1/8} \right\}. \quad (11)$$

The modification induced by the finite size effects SS, Self, QM and DD are all encoded in σ , while the SO contribution is in β .

References

1. B. Abbott et al, Phys. Rev. D **69**, 122001 (2004).
2. E. Messaritaki, Class. Quantum Grav. **22**, S1119 (2005).
3. E. Berti, A. Buonanno, and C. M. Will, Phys. Rev. D **71**, 084025 (2005).
4. L. Á. Gergely, Z. Perjés, and M. Vasúth, Astrophys. J. Suppl. **126**, 79 (2000).
5. L. Á. Gergely, Z. Keresztes, and B. Mikóczi, Astrophys. J. Suppl. **167**, 286 (2006).
6. L. Á. Gergely, Z. I. Perjés, and M. Vasúth, Phys. Rev. D **58**, 124001 (1998).
7. L. Á. Gergely, Phys. Rev. D **61**, 024035 (2000).
8. L. Á. Gergely, Z. Keresztes, Phys. Rev. D **67**, 024020 (2003).
9. M. Vasúth, Z. Keresztes, A. Mihály, and L. Á. Gergely, Phys. Rev. D **68**, 124006 (2003).
10. A. Gopakumar, Bala R. Iyer Phys. Rev. D **56**, 7708 (1997).
11. Z. Keresztes, B. Mikóczi and L. Á. Gergely Phys. Rev. D **72**, 104022 (2005).
12. G. Schäfer, N. Wex, Phys. Lett A **174**, 196 (1993), erratum: **177**, 461 (1993).
13. L. Kidder, Phys. Rev. D **52**, 821 (1995).
14. B. Mikóczi, M. Vasúth, and L. Á. Gergely Phys. Rev. D **71**, 124043 (2005).
15. L. Blanchet, Phys. Rev. D **54**, 1417 (1996).
16. L. Blanchet, G. Faye, B. R. Iyer, and B. Joguet Phys. Rev. D **65**, 061501 (2002); Erratum ibid. D **71**, 129902 (2005).
17. L. Blanchet, A. Buonanno, and G. Faye Phys. Rev. D **74**, 104034 (2006); Erratum ibid. D **75**, 049903 (2007).